GREAT EAST LAKE

2021 SAMPLING HIGHLIGHTS

Station – 1 Center

Acton, ME and Wakefield, NH



Station 1 Center was used as a reference point to represent the overall Great East Lake water quality. Water quality data displayed in Tables 1 and 2 are surface water measurements, with the exception of the dissolved oxygen data that were collected near the lake bottom.

Blue = Oligotrophic

Yellow = Mesotrophic

Red = Eutrophic

Gray = No Data




Figure 1. Great East Lake Water Quality (2021)


Table 1. 2021 Great East Lake Seasonal Averages and NH DES Aquatic Life Nutrient Criteria¹

Parameter	Oligotrophic	Mesotrophic	ophic Eutrophic Great East Lake Average (range)		Great East Lake Classification
Water Clarity (meters)	4.0 – 7.0	2.5 - 4.0	< 2.5	10.4 meters (9.5 – 10.9)	Oligotrophic
Chlorophyll <i>a</i> ¹ (ppb)	< 3.3	> 3.3 – 5.0	> 5.0 – 11.0	1.5 ppb (1.1 – 1.7)	Oligotrophic
Total Phosphorus ¹ (ppb)	< 8.0	> 8.0 – 12.0	> 12.0 – 28.0	3.8 ppb (3.2–4.7)	Oligotrophic
Dissolved Oxygen (mg/L)	5.0 – 7.0	2.0 – 5.0	<2.0	6.0 mg/L (3.5–9.4)	Oligotrophic

^{*} Dissolved oxygen concentrations were measured between 11.5 and 31.0 meters, in the bottom water layer, on August 26, 2021.

Table 2. 2021 Great East Lake Seasonal Average Accessory Water Quality Measurements

Parameter	Assessment Criteria					Great East Lake Average (range)	Great East Lake Classification
Color (color units)	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored	9.1 color units (range: 7.5 – 10.7)	Uncolored
Alkalinity (mg/L)	< 0.0 acidified	0.1 – 2.0 extremely vulnerable	2.1 – 10 moderately vulnerable	10.1 – 25.0 low vulnerability	> 25.0 not vulnerable	8.1 mg/L (range: 7.4 – 8.3)	Moderately vulnerable
pH (std units)	< 5.5 suboptimal for successful growth and reproduction		6.5 – 9.0 optimal range for fish growth and reproduction			6.6 standard units (range: 6.4 – 6.9)	Optimal range for fish growth and reproduction
Specific Conductivity (uS/cm)	< 50 uS/cm Characteristic of minimally impacted NH lakes		50-100 uS/cm Lakes with some human influence	> 100 uS/cm Characteristic of lakes experiencing human disturbances		82.9 <i>u</i> S/cm (range: 82.0 – 84.3)	Lakes with some human influence






Figure 2 and 3. Seasonal Secchi disk transparency, chlorophyll a concentrations and dissolved color concentrations. Figures 2 and 3 illustrate the interplay among Secchi Disk transparency, chlorophyll a and dissolved color. Shallower water transparency measurements oftentimes correspond to increases in chlorophyll a and/or color concentrations.


LONG-TERM TRENDS

WATER CLARITY: The Great East Lake water clarity measurements, measured as Secchi Disk transparency, display a trend of increasing water clarity (Figure 4).

CHLOROPHYLL: The Great East Lake chlorophyll a concentrations, a measure of microscopic plant life within the lake, display a trend of decreasing concentrations (Figure 4).

TOTAL PHOSPHORUS: Phosphorus is the nutrient most responsible for microscopic plant growth. The Great East Lake total phosphorus concentrations have varied among years (Figure 5). The total phosphorus data do not display a long-term water quality trend.

COLOR: The Great East Lake color data, the result of naturally occurring "tea" color substances from the breakdown of soils and plant materials, display a trend of decreasing concentrations (Figure 5).







Table 3. Salmon Falls Headwaters Seasonal Average Water Quality Inter-comparison (2021)

Lake	Average Secchi Disk Transparency (meters)	Average Chlorophyll <i>a</i> (ppb)	Average Total Phosphorus (ppb)	Average Dissolved Oxygen (mg/L)	
Great East Lake	10.4	1.5	3.8	6.0	
Wilson Lake	Not Assessed				
Lovell Lake	7.0	2.9 7.1		1.9	
Horn Pond	Not Assessed				
Lake Ivanhoe 4.2		4.8	8.2		

- Water quality data are reported for a deep reference sampling location in each water body
- Dissolved oxygen measurements were collected in the summer (late July and August) in the bottom water layer (metalimnion or hypolimnion).
- ------ Indicates the site is too shallow to form a bottom water layer (metalimnion or hypolimnion) during the summer months.

Figures 4 and 5. Changes in the Great East Lake water clarity (Secchi Disk depth), chlorophyll a, dissolved color and total phosphorus concentrations measured between 1974 and 2021. These data illustrate the relationship among plant growth, water color and water clarity. Total phosphorus data are also displayed and are oftentimes correlated with the amount of plant growth. Trendlines are displayed when sufficient data are available. Longterm trends are based on the analysis of annual median values

Figure 6. Great East Lake dissolved oxygen profiles collected July 22 through August 26, 2021. The vertical red line indicates the oxygen concentration of five milligrams per liter that is commonly considered the threshold for successful growth and reproduction of cold water fish such as trout and salmon.



Recommendations

Implement Best Management Practices within the Great East Lake watershed to minimize the adverse impacts of polluted runoff and erosion on the lake. Refer to "Landscaping at the Water's Edge: An Ecological Approach" and "New Hampshire Homeowner's Guide to Stormwater Management: Do-It-Yourself Stormwater Solutions for Your Home" for more information on how to reduce nutrient loading caused by overland run-off. The Acton Wakefield Watersheds Alliance also offers technical assistance to help design and implement erosion control projects that protect and improve the water quality.

- https://extension.unh.edu/resources/files/resource004159_rep5940.pdf
- https://www.des.nh.gov/sites/g/files/ehbemt341/files/documents/2020-01/homeowner-guide-stormwater.pdf
- https://awwatersheds.org/conservation-practices-for-homeowners//

Figure 7. Great East Lake Acton, ME and Wakefield, NH

2021 Deep water sampling sites and seasonal average water transaprency.

